


CLAIMS

1. A method of treating a disease, damage or disorder of the central nervous system associated with a disorder of neurochemical equilibrium of a biogenic amine or other neurotransmitter, comprising administering to a subject in need thereof a compound of formula I



wherein


X is selected from the group consisting of CH₂, O, S, S(=O), S(=O)₂ and NR^a,
wherein R^a is selected from the group consisting of hydrogen, C₁-C₃-alkyl, C₁-C₃-alkanoyl, C₁-C₇-alkoxycarbonyl, C₇-C₁₀-arylalkyloxycarbonyl, C₇-C₁₀-aroyl, C₇-C₁₀-arylalkyl, C₃-C₇-alkylsilyl and C₅-C₁₀-alkylsilylalkyloxyalkyl;

Y and Z are each independently selected from the group consisting of hydrogen, halogen, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, halo-C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkanoyl, amino, amino-C₁-C₄-alkyl, C₁-C₄-alkylamino, *N*-(C₁-C₄-alkyl)amino, *N,N*-di(C₁-C₄-alkyl)amino, thiol, C₁-C₄-alkylthio, sulfonyl, C₁-C₄-alkylsulfonyl, sulfinyl, C₁-C₄-alkylsulfinyl, carboxy, C₁-C₄-alkoxycarbonyl, cyano and nitro;

R¹ is selected from the group consisting of hydrogen, halogen, C₁-C₇-alkyl optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl; C₂-C₇-alkenyl optionally substituted with one or more halogen atoms; C₂-C₇-alkynyl; hydroxy; hydroxy-C₂-C₇-alkenyl; hydroxy-C₂-C₇-alkynyl; C₁-C₇-alkoxy; thiol; thio-C₂-C₇-alkenyl; thio-C₂-C₇-alkynyl; C₁-C₇-alkylthio; amino, *N*-(C₁-C₇-alkyl)amino; *N,N*-di(C₁-C₇-alkyl)amino; C₁-C₇-alkylamino; amino-C₂-C₇-alkenyl; amino-C₂-C₇-alkynyl; amino-C₁-C₇-alkoxy; C₁-C₇-alkanoyl; C₇-C₁₀-aroyl; oxo-C₁-C₇-alkyl; C₁-C₇-alkanoyloxy; carboxy; $\#$ C₁-C₇-alkyloxycarbonyl \neq C₁-C₇-aryloxycarbonyl; carbamoyl; *N*-(C₁-C₇-

alkyl)carbamoyl; *N,N*-di(C₁-C₇-alkyl)carbamoyl; cyano; cyano-C₁-C₇-alkyl; sulfonyl; C₁-C₇-alkylsulfonyl; sulfinyl; C₁-C₇-alkylsulfinyl; nitro;

a substituent of the formula II:



II


wherein

R³ and R⁴ are each independently selected from the group consisting of hydrogen, C₁-C₆-alkyl, and aryl; or

R³ and R⁴ taken together with the nitrogen atom to which they are attached form a heterocycle or heteroaryl group optionally substituted with one or two substituents which are selected from the group consisting of halogen, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, and C₁-C₄ alkylsulfinyl;

m and n are each independently an integer from 0 to 3;

Q₁ and Q₂ are each independently selected from the group consisting of oxygen, sulfur,



wherein substituents

y₁ and y₂ are each independently selected from the group consisting of hydrogen, halogen, C₁-C₄-alkyl or aryl optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl; or aryl optionally substituted with one or two substituents selected from the group consisting of halogen, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino,

sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, and C₁-C₄ alkylsulfinyl; hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkanoyl, thiol, C₁-C₄-alkylthio, sulfonyl, C₁-C₄-alkylsulfonyl, sulfinyl, C₁-C₄-alkylsulfinyl, cyano, and nitro, or

y₁ and y₂ taken together with the carbon atom to which they are attached form a carbonyl group or an imino group;


a monocyclic or bicyclic aryl group; a monocyclic or bicyclic heteroaryl group; and a heterocycle, wherein the monocyclic or bicyclic aryl group, the monocyclic or bicyclic heteroaryl group and the heterocycle are linked to the thiophene ring via a direct bond or a C₁-C₄ alkylene group, and are each optionally substituted with one or more substituents selected from the group consisting of fluoro, chloro, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl;

R² is hydrogen, C₁-C₇-alkyl optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl; aryl optionally substituted with one or two substituents selected from the group consisting of halogen, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, and C₁-C₄ alkylsulfinyl; C₁-C₇-alkanoyl, C₁-C₇-alkoxycarbonyl, C₇-C₁₀-arylalkyloxycarbonyl, C₇-C₁₀-aroyle, C₇-C₁₀-arylalkyl, or C₃-C₇-alkylsilyl;

and a pharmaceutically acceptable salt or solvate thereof.

2. The method of claim 1, wherein the biogenic amine is serotonin, norepinephrine or dopamine.
3. The method of claim 1, wherein the neurotransmitter is glutamate.
4. The method of claim 1 wherein the compound of formula I regulates the synthesis, storage, release, metabolism, reabsorption or receptor binding of a biogenic amine or neurotransmitter.
5. The method of claim 4, wherein the compound of formula I binds to a receptor of a biogenic amine.
6. The method of claim 5, wherein the compound of formula I binds to a serotonin 5-HT_{2A} or 5-HT_{2C} receptor.

7. The method of claim 6, wherein the compound of formula I binds to a serotonin 5-HT_{2A} or 5-HT_{2C} receptor with an IC₅₀ of less than 1 μ M.
8. The method of claim 1, wherein the compound of formula I binds to a σ 1 receptor with an IC₅₀ of less than 1 μ M.
9. The method of claim 1, wherein the compound of formula I binds to a σ 1 receptor and to at least one serotonin receptor selected from 5-HT_{2A} and 5-HT_{2C}.
10. The method of claim 1, wherein the disease or disorder of the central nervous system ~~are~~ is selected from the group consisting of anxiety, depression, bipolar disorders, sleeping disorders, sexual disorders, psychosis, borderline psychosis, schizophrenia, migraine, personality disorders, obsessive-compulsive disorders, social phobia, ~~or~~ panic attacks, organic mental disorders in children, aggression, memory disorders, personality disorders in elderly people, addiction, obesity, bulimia and other eating disorders, snoring, and premenstrual troubles.
11. The method of claim 1, wherein the damage to the central nervous system is caused by trauma, brain stroke, neurodegenerative diseases, cardiovascular disorders, thrombosis, infarct or gastrointestinal disorders.
12. The method of claim 1 wherein X is O, S, or NR^a, wherein R^a is hydrogen, C₁-C₃-alkyl, C₁-C₃-alkanoyl, C₇-C₁₀-aroyl or C₇-C₁₀-arylalkyl.
13. The method of claim 1, wherein Y and Z are each independently selected from the group consisting of hydrogen, fluorine, chlorine, bromine, C₁-C₄-alkyl, halo-C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkanoyl, amino, amino-C₁-C₄-alkyl, N-(C₁-C₄-alkyl)amino, N,N-di(C₁-C₄-alkyl)amino, thiol, C₁-C₄-alkylthio, cyano and nitro.
14. The method of claim 1, wherein R¹ is selected from the group consisting of hydrogen, halogen, C₁-C₄-alkyl optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, N-(C₁-C₄) alkylamino and N,N-di(C₁-C₄-alkyl)-amino; hydroxy; C₁-C₄ alkoxy; thiol; C₁-C₄ alkylthio; amino; N-(C₁-C₄ alkyl)amino; N,N-di-(C₁-C₄ alkyl)amino; C₁-C₇ alkanoyl; C₇-C₁₀-aroyl; C₁-C₇ alkanoyloxy; C₁-C₇ alkyloxycarbonyl, C₇-C₁₀-aryloxycarbonyl; carbamoyl; N-(C₁-C₇-alkyl)carbamoyl; N,N-di(C₁-C₇-alkyl)carbamoyl; cyano; cyano-C₁-C₇ alkyl; nitro;
and a substituent of the formula II:



II

wherein

R^3 and R^4 are each independently hydrogen, C_1 - C_4 -alkyl, or aryl, or

R^3 and R^4 taken together with the nitrogen atom to which they are attached form a heterocycle or heteroaryl group selected from the group consisting of morpholine-4-yl, piperidine-1-yl, pyrrolidine-1-yl, imidazole-1-yl and piperazine-1-yl;

m and n are each independently an integer from 0 to 3; and

Q_1 and Q_2 are each independently oxygen or CH_2 .

15. The method of claim 1, wherein the compound of formula I is selected from the group consisting of:

1H-8-oxa-1-aza-dibenzo[e,h]azulene;

11-chloro-1H-8-oxa-1-aza-dibenzo[e,h]azulene;

1H-8-thia-1-aza-dibenzo[e,h]azulene;

1H-8-oxa-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

11-chloro-1H-8-oxa-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

1H-8-thia-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1-aza-dibenzo[e,h]azulene-2-carbaldehyde;

[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-yl]-methanol;

[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-yl]-methanol;

[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1-aza-dibenzo[e,h]azulen-2-yl]-methanol;

dimethyl-[2-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl]-amine;

dimethyl-[2-(1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl]-amine;

dimethyl-3-(1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

{2-[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl}-dimethyl-amine;

[2-(11-chloro-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-dimethyl-amine;

{3-[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl}-dimethyl-amine;

[3-(11-chloro-1H-8-oxa-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-dimethyl-amine;

dimethyl-2-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl]-amine;

dimethyl-2-(1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl]-amine;

dimethyl-3-(1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy]-propylamine;

3-(1H-8-thia-1-aza-dibenzo[e,h]azulen-2-ylmethoxy)-propylamine; and

a pharmaceutically acceptable salt or solvate thereof.